Systematics in Next Generation Oscillation Experiments

Joe Walsh

walshj29@msu.edu

XXIII International Workshop on Next Generation Nucleon Decay and Neutrino Detectors

Friday November 1st 2024

CBPF, Rio de Janeiro









Office of Science

An Age of Precision

Where we are...



...where we want to get




CKM Unitarity Triangles Eur. Phys. J. C 2005, 41, 1–131.


arXiv:2212.00809

The Next Generation















Measuring Oscillations

Not a pure probability measurement, in spectra predictions mixing effect is convolved with flux, interaction, detector modeling

$$R = \Phi \times \sigma \times \epsilon \times P(\nu_{\alpha} \to \nu_{\beta})$$



$$L = \underbrace{ \left[L_{Flux} \, \otimes L_{xsec} \, \otimes L_{Det}
ight] \otimes L_{PMNS} }_{ ext{Near Detector}}$$


Approaches to Systematics

Model Driven

Use data to constrain systematic models – Can help you understand underlying physics

Need adequate model freedoms to describe data

Significant risk of model dependent bias in results




Data Driven

Use data from one detector to more directly make predictions for a functionally identical or similar

Avoids model dependencies

Still need to understand any possible differences between very similar detectors like size, flux, or geometry





General Considerations for Neutrino Experiments

Neutrino Sources: Nuclear

Reactors

Many contributing beta decay processes each with fine structure. Models have significant discrepancy






See PROSPECT Talk from Tuesday [O. B. Rodrigues]

Neutrino Sources: Hadron Decays

Beam and Atmospheric

Hadronic production processes and secondary interactions are biggest contribution to flux uncertainty



J-PARC neutrino beam

For beams, horn current and field have impact too as well as beamline geometry

Hadron production

See Lu Ren's talk from Wednesday




On production


talk from Wednesday

Detailed measurements of hadrons exiting fixed targets in proton beam at different energies

Mix of thin-target and replica-target measurements Reduces T2K flux uncertainty from ~10% to ~5%



EMPHAT C



NuMI Horn 1


EMPHATIC

Also makes hadron measurements

Will make first measurement of hadrons exiting neutrino beam focusing horn


Interaction modeling

Each interaction mode has different relationship between neutrino energy and final-state kinematics




CCQE and Multinucleon Knockout Recent measurements seem to disfavor dipole form factor model

We cannot yet accurately model pion kinematics in resonant interactions



Hadronization models and tools from collider world may not accurately describe relatively low-W systems

Are our corrections sufficient?





A complicated nuclear environment

Interactions often take place in the complicated nuclear environment

Ground state impacts cross section and available energy in non-trivial ways, need to go beyond PWIA

Final State interactions can alter FS-topology and kinematics introducing irreducible backgrounds and smearing energy





Charge Exchange

Can uses TKI to probe nuclear effects and even separate out free proton interactions

Elastic Scattering


Detector Systematics

Need to understand how particles and different interaction topologies look in our detectors

Understand how particles propagate and deposit energy in the detector, and how accurately we can record it electronically

Geometry can play a big roll in near-to-far acceptance differences, as well as reconstruction efficiencies and fiducial volume mass uncertainties

Energy scale is very important. Independent measures as well as calibration sources can be used to assess this



Invariant Mass 2R e-like


ND µ
Shower

ND may not contain all events large FD can

Physical surveys and crossing tracks useful for assessing geometry

Develop control samples for specific systematics

- Stopping particles
- Through-going cosmics
- Standard candles and decays



https://doi.org/10.1016/S0168-9002(03)00425-X

How systematics impact oscillations


Neutrino mode: appearance

Example from Hyper-K

If ν_e and ν_μ energy scales are not similar, and we don't understand the ν_e energy scale well we bias appearance shape

For CP-conserving values of δ_{CP} there are subtle shape changes from subdominant terms in the oscillation

A bias in the energy scale results in inability to exclude false minima in the constraint and determine correct octant



The Next Generation










Reactor, Atmospheric, Solar
Inverse Beta Decay
20 kt Liquid Scintillator

Narrowband Sub-GeV Beam, Atmospheric, Solar Mostly quasielastic 170 kt Water Cherenkov Broadband Multi GeV beam, Atmospheric, Solar Broad mix of QE, RES, SIS, DIS 2-4 x 17 kt LAr-TPC













20 kt LS detector at 52.5 km from two nuclear power plants








 $3\% / \sqrt{E}$ energy resolution 17,612 20" PMTs 25,600 3" PMTs 77% total photocathode coverage


Cosmic ray tracker and outer WC det reduce backgrounds



JUNO-TAO

44 m from reactor
Will measure fine structure in spectrum
with high statistics and
1% uncertainty on spectra at 3 MeV





https://arxiv.org/pdf/2405.18008


Doped with Gd to increase n-capture Helps account for acceptance difference due to smaller size




Lower stats at far detector lead to larger backgrounds


Radiopurity of LS is monitored with dedicated OSIRIS detector

JUNO-OSIRIS





Hyper-Kamiokande






Hyper-K Far Detector

Significant 50cm PMT performance increase from SK Better QE, timing resolution and charge discrimination


Calibration including light injection, radioactive sources and photogrammetry



KM3NET/IceCube style multi-PMTs Better timing and angular resolution.

Better ring separation and lightabsorption constraint





OD PMT+WLS **Underwater Electronics** mPMT 50cm PMT 70cm grid 21



ND280 Upgrade

Hyper-K will inherit the suite of near detectors from T2K Including magnetized off-axis detector ND280

UAI Magnet

ECal

ECal

FCal

HATPC TPC

TPC

TPC

FCD:

uper Fine Grained Detector


Full 3D readout plastic scintillator target detector SFGD


Improved muon acceptance and hadron efficiencies


Double mass of existing targets

Improved electron-photon discrimination

Neutron tagging and kinematics from ToF Allows more TKI variable measurements








T2K is already taking data in upgraded ND280 at over 750 kW

Event number : 345342 | Run number : 16847 | Spill : 28852 | Time : Fri 2024-06-07 18:29:00 JST | Trigger: Beam Spill






Intermediate Water Cherenkov Detector


Hyper-K will supplement the existing suite of T2K near detectors with a Water Cherenkov detector at 830 m capable of sampling different off-axis spectra

Same detection principle and target material as HK FD Benefits from more direct cancellation of systematics


More finely instrumented with mPMT modules

Large size is self-shielding of OOFV γ background to ν_e














Perforated readout strips in Vertical Drift



Wireplane in Horizontal Drift





External optical bench Feed through for UV laser UV laser paths

Optics to reflect laser into cryostat

Drift volume

mirror

DTS time ticks (16ns)

2-4 10 kt FV LAr-TPC detectors


Horizontal Drift: Wire plane readout

Vertical Drift: Copper strip charge amplifier readout planes


26

Photon detection system including Optical X-ARAPUCA detectors provide precise t₀ for events


Laser ionization calibration system for E-field uniformity See talk by José Maneira









Prototype Pixel readout plane







Movable Segmented LAr-TPC with pixel readout

Segmentation leads to smaller drift lengths and better pile-up separation

TMS muon spectrometer catches forward going muons to account for ND-FD acceptance difference and determine wrong-sign contamination

See Zoya Vallari's talk


PRISM

PRISM


See Akutsu-san's talk for more details

Predicting amount of neutrino energy which is misreconstructed is model dependent.

What if we use a linear combination of ND fluxes to create FD flux and minimize interaction model dependence



 $E_{\rm rec} \; (Gev) \\$ Predicted true neutrino energy contributions to reconstructed neutrino energy at Hyper-K far detector




PRISM

See Akutsu-san's talk for more details


Pseudo-monochromatic neutrino spectra

Linear Combination, 0.9 GeV Mean





DUNE PRISM analysis sensitivity with wrong interaction model



Other measurements


Prototypes

Prototypes in test beams help us understand the detectors we are building.




m-PMT design and Hyper-K calibration tested in WCTE

ProtoDUNE and other LAr detectors successfully testing LAr-TPC readout and calibration systems






2024-07-11 19:52:24 UTC



External Measurements


Mix of both neutrino and non-neutrino experiment data can feed into modeling




Healthy cross section community helps keep pushing forward on model development

There are non-neutrino experiments which can tell us about both neutrino interactions and the modeling of final state particles

mini-CAPTAIN LAr-TPC in neutron beam at LANSCE



Electron scattering data can help us learn more about the ground state of the nucleus



Dedicated neutron scattering experiments at relevant energies

MINERvA 3D lepton-proton correlation measurement

Outlook

Outlook

Next generation experiments will be systematically dominated

- ➤ Need robust models and measurements to remain unbiased
- ➤ Near detectors are a powerful tool for characterizing our sources
- ➤ Prototypes help us understand the performance of these technologies
- External measurements by dedicated experiments are valuable
- ➤ Model-independent approaches can reduce our susceptibility to bias

We can do the work now to prepare for the data we will very rapidly collect with these new 10s kt to 100s kt scale detectors and high intensity sources

Obrigado!

Backup

Robustness Studies

"Fake Data Studies"

Test for out-of-model effects which could bias results


Generate "fake data" – a modified set of the nominal MC to include variations not deliberately included inthe uncertainty model

Fit model at ND or use ND tune to predict FD fake data spectra

Check shifts in credible interval positions and sizes are not large relative to systematic uncertainty contribution

Perform this procedure at key several different oscillation parameter combinations of interest

Future experiments may wish to expand this through by testing compatibility with model cloud and not just central values



Example from joint NOvA-T2K analysis based on T2K ND280 CC0 π data


ND-GAr


Magnetized High-Pressure gaseous Argon TPC

Dense enough to collect appreciable statistics


Lower tracking thresholds and more precise momentum and PID

https://arxiv.org/pdf/1910.06422





ND280 Upgrade








Figure 6.21: Comparison of the ability of a SuperFGD and an FGD XY to measure δp_{TT} . The y-axis reports the CCQE-like cross section within the phase space accessible by the relevant detector. Detector smearing and acceptance effects are applied as described in Sec. 6.5.2.

ZK







Improved constraint on Xsec systs




Dashed= pre-upgrade Dotted=Upgrade muon only Solid=Upgrade muon + proton



ND280 now has a much closer match in angular acceptance to WC dets







Impact of pre-upgrade ND280 in T2K analysis on Super-K predictions


Sample		Uncertainty source (%) Flux Interaction FD + SI + PN			Flux⊗Interaction (%)	Total (%)
1Rμ	$\frac{v}{\overline{v}}$	2.9 (5.0) 2.8 (4.7)	3.1 (11.7) 3.0 (10.8)	2.1 (2.7) 1.9 (2.3)	2.2 (12.7) 3.4 (11.8)	3.0 (13.0) 4.0 (12.0)
1Re	$\frac{v}{\overline{v}}$	2.8 (4.8) 2.9 (4.7)	3.2 (12.6) 3.1 (11.1)	3.1 (3.2) 3.9 (4.2)	3.6 (13.5) 4.3 (12.1)	4.7 (13.8) 5.9 (12.7)
1Re1de	ν	2.8 (4.9)	4.2 (12.1)	13.4 (13.4)	5.0 (13.1)	14.3 (18.7

Eur. Phys. J. C 83, 782 (2023)






Calibration methods in WC and LS detectors




Scintillator used to produce photons in SK light-injection systems common in scintillator detectors

From Rhea Gaur Thesis

Photogrammetry can be used to measure detector geometry using underwater survey drones



https://arxiv.org/abs/1307.0162



Cf source coupled with ncapture Ni target Can test prompt+neutron capture coincidence


JUNO calibration system uses fixed and cable movable sources as well as maneuverable vehicles https://arxiv.org/pdf/2011.06405

Free-proton measurements

Exploit FS-kinematic differences between free-proton and nuclear target data

Allows for better understanding of nuclear effects


MINERvA has shown this works, ND280 Upgrade should have even better performance to resolution and timing




https://arxiv.org/pdf/2405.18008

JUNO-only systematics

Systematic effect	Number of parameters	Relative uncertainty [%]		
		input	rate	
JUNO systematic uncertainty	{13+340}		1.4	
$\sin^2 heta_{13}$	1	3.2	0.13	
Matter density (MSW)	1	6.1	0.062	
Detector normalization	1	1.0	1.0	
Energy resolution	3	0.77, 1.6, 3.3	5.4×10^{-7}	
Background rates	$\{7\}$		0.96	
Accidentals	1	1.0	0.019	
$^9{ m Li}/^8{ m He}$	1	20	0.37	
Fast neutrons	1	100	0.23	
$^{13}\mathrm{C}(lpha,n)^{16}\mathrm{O}$	1	50	0.058	
Geoneutrinos	1	30	0.84	
Atmospheric neutrinos	1	50	0.19	
World reactors	1	2.0	0.046	
Background shape	340		0.033	
$^9 \text{Li}/^8 \text{He}$	(340)	10	0.012	
Fast neutrons	(340)	20	0.0026	
$^{13}\mathrm{C}(lpha,n)^{16}\mathrm{O}$	(340)	50	0.0053	
Geoneutrinos	(340)	5.0	0.026	
Atmospheric neutrinos	(340)	50	0.011	
World reactors	(340)	5.0	0.010	
IBD spectrum shape uncertainty from TAO	340	1.3 - 45	0.35	







Systematics in MO sensitivity

Uncertainties	$ \Delta\chi^2_{\rm min} $	$ \Delta\chi^2_{\mathrm{min}} $ change
Statistics of JUNO and TAO	11.5	
+ Common uncertainty	10.8	-0.7
+ TAO uncertainty	10.2	-0.6
+ JUNO geoneutrinos	9.7	-0.5
+ JUNO world reactors	9.4	-0.3
+ JUNO accidental	9.2	-0.2
$+$ JUNO $^9\mathrm{Li}/^8\mathrm{He}$	9.1	-0.1
+ JUNO other backgrounds	9.0	-0.05
Total	9.0	







T2K constraint indicative of power of ND280 for constraining Hyper-K samples.

Upgraded ND280 collects roughly double statistics for same POT with higher acceptance better tracking thresholds

https://arxiv.org/pdf/1805.04163

TABLE IV. Current systematic uncertainty contributions to the T2K oscillation measurements taken from [91] and [92]

$\nu_{\mu} \ CC$	$ u_{ m e}~CC$	$ar{ u_{\mu}} \ CC$	$ar{ u_{ m e}}~CC$
10.8%	10.9%	11.9%	12.4%
2.8%	2.9%	3.3%	3.2%
0.8%	3.0%	0.8%	3.3%
3.9%	2.4%	3.3%	3.1%
1.5%	2.5%	2.1%	2.5%
11.9%	12.2%	13.0%	13.4%
5.1%	5.4%	5.2%	6.2%
	10.8% 2.8% 0.8% 3.9% 1.5%	10.8% 10.9% 2.8% 2.9% 0.8% 3.0% 3.9% 2.4% 1.5% 2.5% 11.9% 12.2%	10.8% 10.9% 11.9% 2.8% 2.9% 3.3% 0.8% 3.0% 0.8% 3.9% 2.4% 3.3% 1.5% 2.5% 2.1% 11.9% 12.2% 13.0%

Right: Expected postfit constraint on systematics relative to priors https://arxiv.org/pdf/2002.03005




Table 4.1: Primary calibration systems and sources that comprise the nominal DUNE FD calibration design along with their primary usage.

System	Primary Usage		
Existing Sources	Broad range of measurements		
μ , predominantly from cosmic ray	Position (partial), angle (partial), electron lifetime, wire response, dE/dx calibration etc.		
Decay electrons, π^0 from beam, cosmic, atm ν	Test of electromagnetic response model		
³⁹ Ar beta decays	electron lifetime (x,y,z,t) , diffusion, wire response		
External Measurements	Tests of detector model, techniques and systems		
ArgoNeuT [86], ICARUS [87, 116, 117], MicroBooNE	Model parameters (e.g., recombination, diffusion)		
DUNE 35 ton prototype [118]	Alignment and t0 techniques		
ArgonTUBE [119], MicroBooNE [120], SBND, ICARUS [121], ProtoDUNE [20]	Test of systems (e.g., Laser)		
ArgoNeuT [122], MicroBooNE [123, 124, 125, 115, 126, 86], ICARUS [127, 128, 129], ProtoDUNE	Test of calibration techniques and detector model (e.g., electron lifetime, Michel electrons, ³⁹ Ar beta decays)		
ProtoDUNE, LArIAT [19], CAP- TAIN [130]	Test of particle response models and fluid flow models		
LArTPC test stands [131, 132, 133, 88]	Light and LAr properties; signal processing techniques		
Monitoring Systems	Operation, Commissioning and Monitoring		
Purity monitors	Electron lifetime		
Photon detection monitoring System	photon detection system (PD system) response		
Thermometers	Temperature, velocity; test of fluid flow model		
Charge injection	Electronics response		
Dedicated Calibration Systems	Targeted (near) independent, precision calibration		
Direct ionization via laser	Position, angle, electric field (x,y,z,t)		
Photoelectric ejection via laser	Position, electric field (partial)		
Neutron injection	Test of SNB signal, neutron capture model		
Proposed Radioactive source deployment	Test of SNB signal model		


Energy scale impact on δ_{CP} measurement in Hyper-K

Electron-neutrino and electronantineutrino candidate predictions for different values of delta.

0 and π have distinguishable spectra, characterized by distortion of the shape induced by $cos\delta$ terms in the appearance probability

Biases in the reconstructed energy of electron neutrinos can skew bias the shape of this sample and preventing the wrong octant of δ from being excluded due to false minima

This is particularly strong if the muonneutrino and electron-neutrino energy scales differ but are assumed to be the same



https://arxiv.org/abs/1412.3086




FIG. 129. Oscillation probabilities as a function of the neutrino energy for $\nu_{\mu} \to \nu_{e}$ (left) and $\bar{\nu}_{\mu} \to \bar{\nu}_{e}$ (right) transitions with L=295 km and $\sin^{2} 2\theta_{13} = 0.1$. Black, red, green, and blue lines correspond to $\delta_{CP} = 0^{\circ}$, 90°, 180° and -90°, respectively. Solid (dashed) line represents the case for a normal (inverted) mass hierarchy.




FIG. 130. Oscillation probabilities of $\nu_{\mu} \to \nu_{e}$ (left) and $\overline{\nu}_{\mu} \to \overline{\nu}_{e}$ (right) as a function of the neutrino energy with a baseline of 295 km. $\sin^{2} 2\theta_{13} = 0.1$, $\delta_{CP} = -90^{\circ}$, and normal hierarchy are assumed. Contribution from each term of the oscillation probability formula is shown separately.





Different contributions from oscillations to Hyper-K electron neutrino appearance

Contributions to DUNE's Flux model

Oscillation analysis uses principle component decomposition for flux uncertainties

Select components are shown along with their most analogous underlying systematics





Eur. Phys. J. C (2020) **80**: 978