A short report of the NNN

Marcio R. Adames¹*

🏛 ¹Federal University of Technology — Paraná, Curitiba, Paraná, Brasil

November 1, 2024

ENSINO PÚBLICO, GRATUITO E DE QUALIDADE



1/50



Outline


2/50

- **Next Generation Nucleon Dacay and Neutrino Detectors**
- Day one
- Day two
- Day three
- Day four
- Day five

Next Generation Nucleon Decay and Neutrino Detectors

Next Generation Nucleon Decay and Neutrino Detectors

M. R. Adames (UTFPR) Summary Talk November 1, 2024



- Story of the origins of the conference.
- The proton decay searches gave us neutrino detectors.

M. R. Adames (UTEPR) Summary Talk November 1, 2024

UIFPR


Day one

Day one



M. R. Adames (UTFPR) Summary Talk November 1, 2024 5/50


T2K latest results and future prospects Jianrun Hu




- New T2K oscillation analysis includes 10% more data in neutrino mode and improved detector systematics.
- CP conservation rejected at 90%.
- First joint oscillation analysis of SK atmospheric + T2K accelerator neutrino has been performed.
- A limited rejection of inverted mass ordering at 90%.
- CP conservation rejected at $\sim 2\sigma$ (depending on the analysis).
- First joint oscillation analysis of NOvA+T2K shows no strong preference on mass ordering.

M. R. Adames (UTEPR) Summary Talk November 1, 2024

New Results from the NOvA experiment & the joint-fit with T2K - Zoya Vallar






NOvA + T2K analysis:


- Excellent precision on Δm_{32}^2 . No strong mass ordering preference.
- $\delta_{CP} = \pi/2$ lies outside 3 σ C.I. for both orderings. If inverted ordering, CP conserving points lie outside 3 σ C.I.

New NOvA analysis:

- Doubled neutrino data set. New samples, simulation and systematics update featured in this analysis.
- Leading single experiment precision on Δm_{32}^2 . 87% preference for Normal Mass Ordering.

Deep Underground Neutrino Experiment: DUNE, Current Status and Perspectives - G. Valdiviesso





DUNE goals:

 Determine the neutrino mass ordering (sign) of Δ_{31}); the octant of θ_{23} (above or below 45°); if CP is violated in the leptonic sector (measure δ_{CP}); if the 3 flavor mixing + MSW model is correct.

Construction update:


- Caverns were inaugurated (huge);
- Second run of ProtoDUNE was completed a few weeks ago.

Brazilian contribution:

- The HD PDS (X-Arapucas).
- Cryogenics for the far detector.


M. R. Adames (UTEPR) Summary Talk November 1, 2024

Status of the Hyper-Kamiokande Experiment Yousuke Kataoka



- HK is the 3^{rd} generation Water Cherenkov Detector in Kamioka (20 times larger statistics than SK).
- HK improve physics sensitivities for various targets.
- HK construction is on-going on schedule and will start operation in 2027.
- Great capabilities for Nucleon Decay Search.

SuperK latest results and future prospects Shintaro Miki



- Atmospheric neutrinos: Analyzed full purewater data set. Prefer normal ordering with $CL_s = 0.077$. SK-Gd data analyses ongoing. Sensitivity to MO is improved by 33%.
- DSNB: Update the flux limit with SK-Gd data. Spectral fit shows 2.3σ excess to null.
- Reactor: First positive observation in SK. correlated with reactor power.
- Nucleon decays: Updates in two modes: $n \to \mu \pi^0$ and $p \to e^+ X$. (No signal excess found)

Underwater and under-ice km-scale detectors: similarities and differences - Tommaso Chiarusi

Neutrino telescopes allow for novel Second-generation detectors are pass-KM3NeT has ORCA and ARCA. studies in high energy Astrophysics ing the discovery stage

Found a very hi energy muon and fundamental Physics.

Baikal-GVD is being con-signal multi-10 PeV muon.


2024 configuration (~0.5 km³)

Ice-Cube produced the first Neutrino structed in lake Baikal/Russia.

Map of the Galaxy.

1450 m

ceCube Neutrino





62 institutes : 22 countries: 5 continents



Astroparticle Research with Cosmics in the Abyss



11 / 50

M. R. Adames (UTFPR) Summary Talk November 1, 2024

JUNO status and future prospects Prof. Xiaonan Li




Although the project delayed, many challenges were overcome and the installation is nearing its end.


Commissioning shows that the performance of (part of) components/detector is better than the requirement.

Expect to start filling phase in December of this year.

Latest results from the Neutrinos-Angra Project Pietro Chimenti








- Neutrinos Angra is successfully monitoring the Angra-II nuclear reactor since 2020.
- The project has been entirely conceived, designed, constructed and run in Brazil. Small local projects are important for brazilian science.
- Neutrino detector are also efficient neutron detector. This can be used to monitor spent fuel movements near the plant.


Clear Excess of reactor ON periods with respect to reactor OFF.

M. R. Adames (UTEPR) Summary Talk November 1, 2024

Weak pion production in neutrino-nucleus scattering. - Alejandro Mariano

 π prod. on N.





- We have calculated the total cross section for weak pionproduction on nucleons.
- We also developed a model to describe quasi elastic (QE) ν and $\bar{\nu}(\sim \text{GeV})$ scattering by nuclei within a nuclear matter quantum hadrodynamics relativistic approach. We introduced the effect of ground-state correlations (GSC) through a momentum distribution calculated perturbatively, and also final-state interactions (FSI) using a perturbed nucleon and Δ propagators dressed by a self-energy with nucleon-nucleon and Δ -nucleon correlations.
- The model takes into account of 2p2h and 3p3h excitations in addition to the CCQE 1p1h ones. Results are promising, and the model worked guit well. It is clear that now to analyze weak pion production (CC π) on the detector nucleus, many additional effects appear regards the production on nucleons.


Day two

Day two



15 / 50 M. R. Adames (UTFPR) Summary Talk November 1, 2024

Overview of neutrino oscillations **David Vanegas Forero**



ALL 3-neutrino osc. parameters (& ordering) are aimed to be measured at future neutrino experiments (given the important improvement in precision).


This is not the full story: Massive neutrinos imply BSM physics

- BSM physics searches already been performed at Osc. Exps.
- With significant improvement in precision. subleading effects could be found for at neutrino oscillation Exps.

PROSPECT-I final oscillation results and

17/50

PROSPECT-II physics goals and detector design - Ohana B. Rodrigues



measurement PRESPECTtrino spectrum

Goals: - Search for sterile neutrino oscillations: - Precise

We have set world-leading limits on sterile neutring oscillations with the final PROSPECT-I dataset

- Neutrino-4 BF ruled out at $> 5\sigma$
- Gallium Anomaly allowed $\Delta m2 < 10eV^2$ rule out at 95% C.L.
- Leads global SBL electron-flavor disappearance limits over most of the 3-10eV2phase space

of PROSPECT-II robustness give us the chance the ^{235}U antineu- to push even further the borders of knowledge we currently have in the field.

Direct Neutrino Mass Measurements Alessandro Schwemmer

KATRIN

- New leading upper limit from KATRIN: $m_{\rm g}$ < 0.45 eV
- Continue measuring, target sensitivity m_e < 0.2 eV
- R&D phase started for atomic source and micro-calorimeters


Project 8


- First frequency-based upper limit from Project 8: m_e< 155 eV
- Development of large-volume CRES and atomic source

ECHo and HOLMES


- New / updated results: $\rm m_{g}{<}$ 19 eV (ECHo) and $\rm m_{g}{<}$ 28 eV (HOLMES)
- Increase number of detectors and activity









18 / 50





M. R. Adames (UTEPR) Summary Talk November 1, 2024

Neutrinoless double beta decay: status and prospects of searches with liquid scintillator detectors - José Maneira



experiment

Neutrinoless double beta decay (only for Majorana neutrinos) a very high priority search:

Complementary to CP violation searches in neutrino oscillations

Liquid scintillator technique has many advantages:

 High mass, low backgrounds, scalability, target-out measurements

SNO+ is already taking high quality liquid scintillator data.

The precise inverse beta decay Giulia Ricciardi

Inverse beta decay (IBD)

$$\bar{\nu}_e + p \rightarrow e^+ + n$$

by far the most important mechanism of interaction of low-energy antineutrinos




precision of theoretical cross section crucial for high statistics experiments

GR, Vissani & Vignaroli JHEP 08 (2022) 212

The new calculations improved the determination of the cross section and assessment of uncertainty! (in comparisson to the 2003 commonly used calculations)

M. R. Adames (UTFPR) Summary Talk November 1, 2024 20 / 50


Application of NuGraph to Proton Decay for **Event Classification ... in DUNE - Tyler Stokes**



- NuGraph is a Graphical Neural Network originally designed for semantically classifying detector hits:
- The network has been expanded to perform event classification for proton decay:
- Iteration over nuclear model configurations can provide a conservative estimate of nuclear model uncertainty:
- NuGraph performance is very encouraging! (0.98 Area Under the Curve).



Search for proton decay via $p \to e^+ \eta$ and $p \to \mu^+ \eta^-$ in Super-Kamiokande - Natsumi Taniuchi



Why eta? $p \rightarrow e^+ \eta$ and $p \rightarrow \mu^+ \eta$


- Similar to golden $p o e^+ \pi_0$: 3 e-like rings
- High efficiency: all decay products are detectable
- Back-to-back emission: monochromatic momentum of η (~310 MeV/c)

Number of candidates consistent with background prediction.

No indication of proton decay. New lifetime estimates.

M. R. Adames (UTFPR) Summary Talk November 1, 2024

Light and Charge yield study in LAr for Sub-keV energies up to 1 MeV - Youssef Sarkis Mobarak



- We present a first principles approach study based on integral equation for interactions in pure LAr.
- We give a physical interpretation of ε_e that allow to describe the N_{ex}/N_i ratio as function of energy and field.
- With just one parameter, $C_{LAr}=0.37\pm0.07(V/cm)^{\delta_L Ar}$ we can describe data for LAr and LXe. Usufull for phenomenology studies.
- The model for Charge yield have a better match with data than NEST, thanks to Bates-Griffing process.
- The publication with a software to produce outputs soon.

M. R. Adames (UTFPR) Summary Talk November 1, 2024

Day three

Day three




24 / 50 M. R. Adames (UTFPR) Summary Talk November 1, 2024

Day three

25 / 50

MicroBooNE Latest Results and Future Prospects Christopher Thorpe





MicroBooNE is LArTPC with a wide ranging physics program covering oscillations, BSM, neutrino-nucleus cross sections, reconstruction, and LArTPC detector physics. We have:

- a number of results probing the MiniBooNE and LSND anomalies (disfavour ν_e expl.).
- produced a number of results searching for BSM particles produced in our neutrino beams (exclude new regions).
- may infer protons in tracks.


ICARUS at the Short-Baseline Neutrino programfirst results - Laura Pasqualini





SBN sensitivity to Sterile Neutrino

Combined analysis of events collected by ICARUS (far detector) and SBND (near detector) in 3 years (6.6x10²⁰ pot) allows:

- 5σ coverage of the parameter area relevant to LSND anomaly
- probe the parameter area relevant to reactor and gallium anomalies







- ICARUS is smoothly running in Physics mode since June 2022, exposed to BNB and NuMI beams.
- The detector response is calibrated with cosmic muons and protons from neutrino interactions, TPC signals and main detector parameters have been accurately characterized and modeled in the simulation.
- Before the start of the joint operation within SBN, ICARUS only is carrying on several analysis.

Status of the Short-Baseline Near Detector at Fermilab - Gustavo Valdiviesso



SBND Timeline



September 2022



December 2022



April 2023



March 2023 LAr filling completed


SBND (and SBN) Science Goals: Sensitive search for sterile neutrino through oscillations; Precision neutrino-nucleus interactions studies; Beyond Standard Model physics searches.

- Data taking began early in July 2024.
- Physics SBND: Sterile Neutrino Searches; Neutrino-Argon Interactions; BSM physics searches.
- World leading statistics for many processes (in three years).
- X-Arapucas Specialized photon sensors developed in Brazil.

tt Generation Nucleon Dacay and Neutrino Detectors Day one Day two Day three

Hadron Production Experiments Lu Ren

How to Make a Neutrino Beam





Experiment to Measure the Production of Hadrons At Testbeam In Chicagoland

Located at Fermilab Test Beam Facility (FTBF)

NA61/SHINE

The SPS Heavy Ion and Neutrino Experiment

Located at CERN SPS North Area

- Measurements of hadron production are critical to constraining the neutrino flux;
- EMPHATIC and NA61/SHINE are two complementary hadron production experiments;
- Both aim to improve flux predictions for current and future: Accelerator-based neutrino experiments, Atmospheric neutrino experiments and Spallation neutron source neutrino experiments

M. R. Adames (UTFPR) Summary Talk November 1, 2024



28 / 50

Day four




Day four




M. R. Adames (UTFPR) Summary Talk November 1, 2024 29 / 50

Next Generation Nucleon Dacay and Neutrino Detectors Day one Day two Day three Day four

Short-Baseline Anomalies and Sterile Neutrinos: where we stand and where we're going - Mark Ross-Lonergan





- The vanilla 3+1 sterile neutrino interpretation of the anomalies is very much in trouble;
- Regardless of your belief in an eV scale sterile neutrino, the anomalies have survived scrutiny and still need explanations!
- Current and next-gen SBL neutrino experiments acting as portal to the dark sector is a thriving direction, motivated at first by anomalies but now as a field in their own right!


30 / 50

M. R. Adames (UTFPR) Summary Talk November 1, 2024

Next Generation Nucleon Dacay and Neutrino Detectors Day one

Near Detectors that Move and Beam Monitors Dr Ryosuke Akutsu

Role of near detectors: near-to-far extrapolation



- Talked about the importance of determining (true) E_{ν} and how current and future experiments can do it.
- Current experiments: fixed near detectors can constrain flux & cross section with required precisions.
- Next generation experiments: Movable detectors need to be able to measure true E_{ν} dependence of cross sections; directly measure (anti-)electron neutrino cross sections.

ND cannot directly constrain Φ(E_v) x σ(E_v) because:

S.Dolan, NOW2022

- True E. is unknown

- All particles associated with interactions cannot be detected


M. R. Adames (UTFPR) Summary Talk November 1, 2024

31/50

LAr purification for LBNF-DUNE: Brazil in-kind contribution - Pascoal Jose Giglio Pagliuso



32/50



- Alternative Innovative filter media for LAr purification were proposed, produced and tested.
- The media were compared the recommended LAr filter used as reference.
- The innovative capacity of the Li-FAU to capture N2 and the higher performance of our innovative media CuMgAI-LDH in O2 capturing lead to two patents.
- The Conceptual Project Design for the Phase 2 Deliverables to LBNF DUNE was completed and is to be implemented in the FDs (45MUS\$).


LAr Purification for LBNF-DUNE Fluid Dynamics Analysis of LAr Purification... - Prof. Dirceu Norlier





- Contaminants $(N_2, O_2 \text{ and } H_2O)$ in argon significantly affect experimental results
- Argon is purified using a filtering media (which captures O_2 and H_2O).
- Typical filters can't remove N_2 , it must be kept low in the argon purchasing specification.
- ullet An effective removal of N_2 impurities during liquid argon circulation was not known and it is desirable.
- Developed a Li-FAU zeolite media that can filter the three contaminants.
- Developed detailed models and simulations for the purification system.
- The proposed model was able to match the experimental results obtained with the ICEBERG cryostat.

The ionisation laser calibration system for DUNE - José Maneira




- DUNE measurements rely on a very good energy performance;
- Calibrations are important to measure model parameters and check reconstruction performance.
- Calibrations will make use of several natural and deployed sources;
- DUNE has proposed a series of improvements over the MicroBooNE laser system, that are being tested at ProtoDUNE;
- Construction, installation and commissioning of PD-HD system completed at CERN. Taking data now!
- Next: PD-VD. And then... far detector construction!

UT CONTROL TO

35 / 50

Novel LArTPC technologies for near detectors Zoya Vallari




We are excited to embrace the challenge of "toomany-neutrinos" in the precision era of neutrino physics. We aim to capture every neutrino interaction—pixel by pixel!! To cope with the extremely high-multiplicity environment of the most powerful accelerator neutrino beam, DUNE ND-LAr employs multiple advanced technologies for next-generation LArTPCs:

- \sim 14M pixels across a 5 imes 7 array of TPC modules.
- Central cathode.
- Two anode planes of PCB tiles with goldplated pixel pads and LArPix ASICs on opposite sides.
- Optically segmented modular design with 50 cm drift.

M. R. Adames (UTFPR) Summary Talk November 1, 2024

DUNE Far Detector: current and new technologies - Andre F. Steklain Lisboa





- Far site excavation is complete
- Next: Building & Site Infrastructure work until mid-2025
- Cryostat warm structure is on its way to US from CERN to be installed in 2025-26
- Far Detector installation in 2026-27
- Purge and fill with argon in 2028
- Physics in 2028 or early 2029
- Beam physics with Near

DUNE is a long-baseline oscillation experiment and neutrino observatory:

- Unique and complementary reach in oscillations, MeVscale neutrinos, and BSM searches.
- DUNE Far Detector is essential to achieve its scientific goals.
- See also other Talks and Posters from DUNE!

M. R. Adames (UTFPR) Summary Talk November 1, 2024

36 / 50

xt Generation Nucleon Dacay and Neutrino Detectors Day one Day two Day three Day four Disconding Company Open Day two Day three Day four Disconding Company Open Day two Day four Disconding Company Open Day four Disconding Company Open Day four Da

COHERENT: Latest Results and Future Prospects Jonathan Link






- CEvNS is the process of a neutrino scattering off of a nucleus without changing its internal structure.
- CEvNS is a Standard Model prediction, it should in fact exist and it should be tested for.
- Coherent is a suit of detectors located at the Spallation Neutron Source at Oak Ridge National Laboratory.
- The main aim of the COHERENT collaboration is to make direct measurements of CEvNS.
- CEvNS were detected in Cesium Iodide detectors (updated in 2022). Also detected by other detectors.

ext Generation Nucleon Dacay and Neutrino Detectors Day one

Coherent Neutrino-Nucleus Scattering Experiment - CONNIE - Carla Bonifazi





Angra - Brazil

- Skipper-CCDs showed to be very promising for detecting lowenergy processes
- Excellent performance in 2021-2023 with flat background and 15 eV energy- threshold
- New CEvNS limit with 18.4 g-days is comparable to previous with larger exposure.
- New competitive limits on vector mediator, DM modulation and millicharged particles.
- The experiment started its next phase with a 16-sensor Multi-Chip-Module.

M. R. Adames (UTFPR) Summary Talk November 1, 2024 38 / 50

t Generation Nucleon Dacay and Neutrino Detectors Day one Day two Day three Day four

Novel Scintillation Detector - Stefan Schoppmann




39 / 50



NuDoubt**: Search For Double Beta Plus Decays





ext Concration Nucleon Dacay and Neutrino Detectors Day one Day two Day three Day four Day fo

The Eos experiment - Lu Ren

timina







- Eos is a performance demonstrator of the brid neutrino detection concept. (Cherenkov and scintillation light)
- Installation finished in March and commissioning completed in May 2024.
- Just completed water-phase calibration datataking and filling of WbLS (Water based Liquid Scintillator) is in progress.
- Will provide valuable input to future kilo-ton scale hybrid detectors, such as Theia.

xt Generation Nucleon Dacay and Neutrino Detectors Day one Day three Day four Day five

The future of neutrino astronomy with TAMBO Pavel Zhelnin

HEPR

Science goals of TAMBO


Discover new ν point sources

Constrain the astro flavor ratio by measuring ν_z component

Characterize the 1-100 PeV

1x 1500

modules



What's next for TAMBO

- Release full simulation ~ 99%
- Hardware development
 - Scintillator construction
 - DAQ system


module

2024

- · Deployment to test sites (US, Peru)
- Interested in joining? Contact (me or Carlos Argüelles) at pzhelnin@g,harvard.edu or carguelles@g,harvard.edu

2026





41 / 50

Day five



Day five

xt Generation Nucleon Dacay and Neutrino Detectors Day one Day three Day four Day four

Photon detection in neutrino physics Ettore Segreto





196 small X-ARAPUCAs installed in the SBND detector at



- Light detection plays a central role in neutrino physics.
- PMT still widely used by the community.
- New technologies are emerging and are being considered/used by some of the next generation experiments.
- New developments and new ideas need to be pursued and encouraged.
- New sensors and detection technique will enable the discoveries that are in front of us.

ANNIE - The Accelerator Neutrino Neutron Interaction Experiment - Matt Wetstein






- ANNIE has achieved many important technical firsts: First detection of neutrinos with Gd-water (beam?); First detection of neutrinos with an LAPPD; First detection of neutrinos with WbLS.
- Gained considerable operational experience w/ these technologies.
- Made significant progress on the reconstruction tools.
- First data with LAPPDs shows powerful imaging capabilities.
- ANNIE is now in a unique position to measure neutrino-nuclear cross sections in water complementary and synergistic the the LAr short baseline program at FNAL.

Next Generation Nucleon Dacay and Neutrino Detectors Day one Day two Day three Day four Day one Day two Day four Day four

Systematics in Next Generation Oscillation Experiments - Joe Walsh





Next generation experiments will be systematically dominated

- Need robust models and measurements to remain unbiased.
- Near detectors are a powerful tool for characterizing our sources.
- Prototypes help us understand the performance of these technologies.
- External measurements by dedicated experiments are valuable.
- Model-independent approaches can reduce our susceptibility to bias.

We can do the work now to prepare for the data we will very rapidly collect with these new 10s kt to 100s kt scale detectors and high intensity sources.

M. R. Adames (UTFPR) Summary Talk November 1, 2024 45 / 50

oct Generation Nucleon Dacay and Neutrino Detectors Day five Day two Day three Day four Day five

Scintillator detectors: SuperFGD, MINERvA, LiquidO - Hikaru Tanigawa



46 / 50

Scintillation detectors









| | MINERVA | SuperFGD | LiquidO |
|------------------------------|--------------------------|-----------------------------|---|
| technology (segmentation) | Scintillator strip | Scintillator cube | Opaque liquid scintillator |
| timeline | Data taking in 2009-2019 | Data taking started in 2023 | R&D since 2013, First experiment (CLOUD) approved |
| energy scale | multi-GeV acc. neutrino | GeV acc. neutrino | multi-purpose MeV-GeV |

Reviewed 3 types of scintillator detectors.

- MINERvA: Scintillator strips, 8t (inner active region); Data taking completed, performance is very well understood; Actively publishing interesting physics results.
- SuperFGD (T2K Near Detector): Scintillator cubes, 2t; High granularity $\rightarrow 4\pi$ acceptance, low threshold; High readout channel density; Just started physics data taking, working on performance studies.
- LiquidO: Opaque liquid scintillator; Light ball formation → self-segmented detector; MeV-GeV application; Good scalability → large scale reactor neutrino experiments CLOUD (5-10t), SuperChooz (O(10kt)).

SuperFGD "A high-granularity detector for improved neutrino interaction measurements in T2K" - Hokuto Kobayashi






cubes $(2 \times 2 \times 2)$

- The SuperFGD consists of scintillation cubes and readout lights from 3-directions. (Gives short tracks detection and 4π acceptance).
- We constructed and commissioned the SuperFGD. (Cube & box \rightarrow Fiber & MPPC \rightarrow Calibration system \rightarrow Cable \rightarrow "Commissioning with electronics" \rightarrow "operation")
- Initial detector calibration and evaluation were performed. (Gain(\sim 15 ADC/p.e.) and light yield(for MIP, \sim 45 p.e.) are obtained. Further understanding of the detector performances is in progress.)
- From June 2024, we started stable operation to take neutrino beam data with the fully upgraded ND280. (Physics analysis with upgraded ND280 is on going.)



ext Generation Nucleon Dacay and Neutrino Detectors Day one Day three Day four Day force

Constraining the (3+2) sterile neutrino scenario with the IceCube experiment. - Alexander Argüello Quiroga



| | $\sin^2 2\theta_{24}$ | | $\Delta m_{41}^{2} \; [eV^{2}]$ | $\Delta m_{51}^{2} \; [\text{eV}^{2}]$ | $\Delta \chi^2$ |
|----------------------------------|-----------------------|----------------------|---------------------------------|--|-----------------|
| Case (A) from Ref. [35] | | | | 13.9 | 4.32 |
| Case (B) from Ref. [36] | 9.1×10^{-2} | 6.8×10^{-2} | 0.46 | 0.77 | 36.43 |

- Anomalies in the SBL, reactor and Solar experiments could indicate the presence of sterile(s) neutrinos;
- In principle the number of sterile neutrinos is free;
- To IceCube, more sterile states means more resonances that also means stronger contraints;
- Only 4 parameter are needed to get a conservative constraint;
- We performed a binned Poisson likelihood analysis with varius systematics and statistical uncertanties;
- The bounds were compared with the allowed regions of the SBL (A) and (B)
- Case (A) in compatible with IceCube data; Case (B) seems in tension

48 / 50

ext Generation Nucleon Dacay and Neutrino Detectors Day one Day two Day three Day four Day five

Neutrino decoherence in Normal matter Prof. Sarira Sahu



49 / 50

- In this work we have considered the effects of the non-forward neutrino scattering processes on the propagation of neutrinos in a normal matter background.
- We calculated the contribution to the imaginary part of the neutrino thermal self-energy (2-loops) from the non-forward neutrino scattering processes.
- A precise prescription is given to determine the jump operators.

Thank you all for the great workshop!!!

50 / 50